

NEW APPLICATIONS OF THE BIEDENHARN-TEMPLE OPERATOR ⁽¹⁾

P. A. HORVÁTHY

Département de Mathématiques, Université,
Parc de Grandmont, F-37200 TOURS (France)
e-mail: horvathy@univ-tours.fr

Abstract.

The Biedenharn approach to the Dirac-Coulomb problem is applied to a system considered by D'Hoker and Vinet, which consists of a spin $\frac{1}{2}$ particle in the combined field of a Dirac monopole plus a λ^2/r^2 potential. The explicit solution is obtained by diagonalizing the 'Biedenharn-Temple operator', Γ .

1. Introduction.

In a paper anticipating supersymmetric quantum mechanics, [1], Biedenharn proposed a new approach to the Dirac-Coulomb problem. He starts with the square of the Dirac equation. This equation can be written in a non-relativistic Coulomb form (but with a fractional angular momentum), by using the 'Biedenharn-Temple' [1, 2] operator Γ . The solutions of the first-order equation can be recovered from those of the second-order equation by projection.

Recently [3], we have applied Biedenharn's method to the 'dyon' system of D'Hoker and Vinet [4]. Here we illustrate the Biedenharn method on yet another example (also considered by D'Hoker and Vinet [5]), namely that of a spin $\frac{1}{2}$ particle described by the four-component Hamiltonian

$$(1.1) \quad H = \begin{pmatrix} H_1 & \\ & H_0 \end{pmatrix} = \frac{1}{2} \left\{ \pi^2 - q \frac{\boldsymbol{\sigma} \cdot \hat{\mathbf{r}}}{r^2} + \frac{\lambda^2}{r^2} - \lambda \gamma^5 \frac{\boldsymbol{\sigma} \cdot \hat{\mathbf{r}}}{r^2} \right\}$$

where λ is a real constant.

2. The Biedenharn-Temple operator for a monopole system.

The Hamiltonian (1.1) can be viewed as associated to a static gauge field on $\mathbf{R}^4$, $\mathbf{A} = q\mathbf{A}_D$, $A_4 = \lambda/r$, where $\mathbf{A}_D$ denotes the vector potential of a Dirac monopole of unit strength. The square of the associated Dirac operator,

$$(2.1) \quad \not{D} := \begin{pmatrix} & Q^\dagger \\ Q & \end{pmatrix} = \begin{pmatrix} & \boldsymbol{\sigma} \cdot \boldsymbol{\pi} - i\frac{\lambda}{r} \\ \boldsymbol{\sigma} \cdot \boldsymbol{\pi} + i\frac{\lambda}{r} & \end{pmatrix},$$

is seen to be (1.1). Since the Dirac operator is chiral-supersymmetric in any even dimension, the partner hamiltonians have the same spectra.

The conserved total angular momentum is $\mathbf{J} = \mathbf{L} + \boldsymbol{\sigma}/2$, where $\mathbf{L} = \boldsymbol{\ell} - q\hat{\mathbf{r}}$, $\boldsymbol{\ell} = \mathbf{r} \times \boldsymbol{\pi}$. The eigenvalues of $\mathbf{J}^2$ are $j(j+1)$, $j = q - 1/2, q + 1/2, \dots$. Set

$$(2.2) \quad w = \boldsymbol{\sigma} \cdot \hat{\mathbf{r}}, \quad z = \boldsymbol{\sigma} \cdot \boldsymbol{\ell} + 1 \quad \text{and} \quad \mathcal{K} = - \begin{pmatrix} & iz \\ iz & \end{pmatrix}.$$

Note that $w^2 = 1$, and that z anticommutes with w and $\boldsymbol{\sigma} \cdot \boldsymbol{\pi}$,

$$(2.3) \quad \{z, w\} = 0 \quad \text{and} \quad \{z, \boldsymbol{\sigma} \cdot \boldsymbol{\pi}\} = 0.$$

⁽¹⁾ Dedicated to Prof. L. C. Biedenharn, on his 70th birthday. *Festschrift in honor of L. C. Biedenharn*, ed. B. Gruber. Plenum : New York (1994).

$\mathcal{K}$ commutes with the Dirac operator $\not{D}$. Using $(\boldsymbol{\sigma} \cdot \mathbf{L})^2 = \mathbf{L}^2 + i\boldsymbol{\sigma}(\mathbf{L} \times \mathbf{L}) = \mathbf{L}^2 - \boldsymbol{\sigma} \cdot \mathbf{L}$, one proves that $\mathcal{K}^2 = z^2 = \mathbf{J}^2 + 1/4 - q^2$. Thus, z (and $\mathcal{K}$) have *irrational* eigenvalues,

$$(2.4) \quad \kappa = \pm \sqrt{(j + 1/2)^2 - q^2}.$$

Since $j \geq q - 1/2$, $\mathcal{K}$ is hermitian, but for $j = q - 1/2$ its eigenvalue κ vanishes and $\mathcal{K}$ is not invertible.

Using $\boldsymbol{\sigma} \cdot \boldsymbol{\pi} = -iw(\partial_r + 1/r - z/r)$, the supercharges Q and $Q^\dagger$ can be written as

$$(2.5) \quad \begin{cases} Q = -iw\left(\partial_r + \frac{1}{r} - \frac{y}{r}\right) = -i\left(\partial_r + \frac{1}{r} + \frac{x}{r}\right)w, \\ Q^\dagger = -iw\left(\partial_r + \frac{1}{r} - \frac{x}{r}\right) = -i\left(\partial_r + \frac{1}{r} + \frac{y}{r}\right)w, \end{cases}$$

where we introduced the notations

$$(2.6) \quad x = z - \lambda w \quad \text{and} \quad y = z + \lambda w.$$

($xw = -wy$). Let us define the *Biedenharn-Temple operator* as

$$(2.7) \quad \Gamma = -(\boldsymbol{\sigma} \cdot \boldsymbol{\ell} + 1 + \gamma^5 \lambda w) \quad \text{i.e.} \quad -(z + \gamma^5 \lambda w) \equiv -\begin{pmatrix} y \\ x \end{pmatrix}.$$

Since $\Gamma^2 = z^2 + \lambda^2 = \mathbf{J}^2 + 1/4 + \lambda^2 - q^2$, the eigenvalues of Γ are

$$(2.8) \quad \gamma = \pm \sqrt{(j + 1/2)^2 + \lambda^2 - q^2}, \quad \text{sign } \gamma = \text{sign } \kappa.$$

In terms of Γ , $\not{D}^2$ becomes

$$(2.9) \quad \not{D}^2 = \begin{pmatrix} Q^\dagger Q & \\ & Q Q^\dagger \end{pmatrix} = -(\partial_r + \frac{1}{r})^2 + \frac{\Gamma(\Gamma + 1)}{r^2}.$$

Although Γ does not commute with $\not{D}$, it is conserved for the quadratic dynamics.

3. The explicit solution.

The operator Γ can be diagonalized generalizing the procedure in Ref. 6. Let us first assume that $j \geq j + 1/2$, and let $L_\pm := j \pm 1/2$ be the orbital angular momentum quantum number. Consider first the two-component spinors

$$(3.1) \quad \varphi_\pm^\mu = \sqrt{\frac{L_\pm + 1/2 \pm \mu}{2L_\pm + 1}} Y_{L_\pm}^{\mu-1/2} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \pm \sqrt{\frac{L_\pm + 1/2 \mp \mu}{2L_\pm + 1}} Y_{L_\pm}^{\mu+1/2} \begin{pmatrix} 0 \\ 1 \end{pmatrix},$$

where the Y 's are the monopole harmonics and the sign $\pm$ refers to the sign of γ . The φ 's satisfy $\mathbf{J}^2 = j(j+1)$, $J_3 = \mu$, ($\mu = -j, \dots, j$), $\mathbf{L}^2 = L_\pm(L_\pm + 1)$. The 2-spinors

$$(3.2) \quad \begin{cases} \chi_+ = \frac{1}{2j+1} \begin{pmatrix} \varphi_+ & + \frac{q}{j+1/2+|\kappa|} \varphi_- \end{pmatrix} \\ \chi_- = \frac{1}{2j+1} \begin{pmatrix} -\frac{q}{j+1/2+|\kappa|} \varphi_+ & + \varphi_- \end{pmatrix} \end{cases}$$

satisfy the important relations

$$(3.3) \quad z \chi_\pm = \pm |\kappa| \chi_\pm \quad \text{and} \quad w \chi_\pm = \chi_\mp,$$

as well as $\mathbf{J}^2 = j(j+1)$, $J_3 = \mu$, ($\mu = -j, \dots, j$). Hence

$$(3.4) \quad \begin{cases} \phi_+ = (|\kappa| + j + \frac{1}{2}) \chi_+ - \lambda \chi_-, & \phi_- = \lambda \chi_+ + (|\kappa| + j + \frac{1}{2}) \chi_- \\ \Phi_+ = (|\kappa| + j + \frac{1}{2}) \chi_+ + \lambda \chi_-, & \Phi_- = -\lambda \chi_+ + (|\kappa| + j + \frac{1}{2}) \chi_- \end{cases}$$

diagonalize x and y ,

$$(3.5) \quad x\phi_{\pm}^{\mu} = \mp |\gamma| \phi_{\pm}^{\mu} \quad \text{and} \quad y\Phi_{\pm}^{\mu} = \mp |\gamma| \Phi_{\pm}^{\mu}.$$

The operator $w = \boldsymbol{\sigma} \cdot \hat{\mathbf{r}}$ interchanges the x and y eigenspinors,

$$(3.6) \quad w\phi_{\pm}^{\mu} = \Phi_{\mp}^{\mu}.$$

For $j = q - 1/2$, no φ_{-} is available and χ_{-} is hence missing. χ_{+} is proportional to the lowest φ_{+} in (3.1). Thus, there are no ϕ_{-} -states in the $\gamma^5 = -1$ sector and no Φ_{+} states in the $\gamma_5 = 1$ sector. However, in each γ^5 sector, (3.2) yields $(2q)$ states, namely

$$(3.7) \quad \phi_{+}^{\mu 0} = \Phi_{-}^{\mu 0} \propto \sqrt{\frac{q+1/2 \pm \mu}{2q+1}} Y_q^{\mu-1/2} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \pm \sqrt{\frac{q+1/2 \mp \mu}{2q+1}} Y_q^{\mu+1/2} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

They are $(+1)$ -eigenstates of w .

The eigenfunctions of $\mathcal{D}^2$ are then found as

$$(3.8) \quad \begin{cases} \Psi_{\pm|\gamma|} = u_{\pm} \begin{pmatrix} \Phi_{\pm} \\ 0 \end{pmatrix} \\ \psi_{\pm|\gamma|} = u_{\pm} \begin{pmatrix} 0 \\ \phi_{\pm} \end{pmatrix} \end{cases} \quad \text{for} \quad \begin{cases} \gamma^5 = 1 \\ \gamma^5 = -1 \end{cases} \quad \text{for } j \geq q + 1/2$$

$$\begin{cases} \Psi_{-}^0 = u_{-}^0 \begin{pmatrix} \Phi_{-} \\ 0 \end{pmatrix} \\ \psi_{+}^0 = u_{+}^0 \begin{pmatrix} 0 \\ \phi_{+} \end{pmatrix} \end{cases} \quad \text{for} \quad \begin{cases} \gamma^5 = 1 \\ \gamma^5 = -1 \end{cases} \quad \text{for } j = q - 1/2$$

Thus, the radial functions $u_{\pm}(r)$ solve the equations

$$(3.9) \quad \left[-(\partial_r + \frac{1}{r})^2 + \frac{\gamma(\gamma+1)}{r^2} - 2E \right] u_{\pm} = 0.$$

This is the wave equation for a free particle except for the fractional ‘angular momentum’ γ . Its solutions is hence given by the Bessel functions,

$$(3.10) \quad u_{\pm}(r) \propto r^{-1/2} J_{|\gamma| \mp \frac{1}{2}}(\sqrt{2E} r).$$

For $\lambda = 0$ we recover the formulae in [6]. The well-known self-adjointness problem in the $j = q - 1/2$ sector shows up in that the eigenvalue γ in Eq. (2.8) vanishes in this case. (Self-adjointness of $\mathcal{D}^2$ requires in fact $|\lambda| \geq 3/2$ [5]).

Another particular value is $\lambda = \pm q$, when the Biedenharn-Temple operator has half-integer eigenvalues,

$$(3.11) \quad \gamma = \pm(j + \frac{1}{2}).$$

In this case, $\gamma(\gamma+1)$ is the same for $-|\gamma|$ as for $|\gamma| - 1$, leading to identical solutions for (3.9). Thus, the corresponding energy levels are two-fold degenerate. (This only happens for $|\gamma| \geq |\gamma|_{\min} + 1$ i.e. for $j \geq q + 1/2$).

This can also be understood by noting that, for $\lambda = \pm q$, the spin dependence drops out in one of the γ^5 -sectors. For $\lambda = q$, e.g., the Hamiltonian (1.1) reduces to

$$(3.12) \quad H = \begin{pmatrix} H_1 & \\ & H_0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \pi^2 - 2q \frac{\boldsymbol{\sigma} \cdot \hat{\mathbf{r}}}{r^2} + \frac{q^2}{r^2} & \\ & \pi^2 + \frac{q^2}{r^2} \end{pmatrix},$$

i.e. H_0 describes a spin 0 particle, while H_1 corresponds to a particle with anomalous gyromagnetic ratio 4. The system admits hence an extra $o(3)$ symmetry, generated by the spin vectors

$$(3.13) \quad \begin{array}{ll} \mathbf{S}_0 = \frac{1}{2}\boldsymbol{\sigma} & \mathbf{S}_1 = U^\dagger \mathbf{S}_0 U \\ \text{for } H_0 & \text{for } H_1 \end{array},$$

where $U = Q/\sqrt{H_1}$ and $U^{-1} = U^\dagger = 1/\sqrt{H_1} Q^\dagger$ are the unitary transformations which intertwine the non-zero-energy parts of the chiral sectors.

Each of the partner Hamiltonians H_1 and H_0 in (1.1) have a non-relativistic conformal $o(2, 1)$ symmetry [7] which combines, together with $\not{D}$ and $-i\gamma^5 \not{D}$, into an $osp(2, 1)$ superalgebra [5]. Since the Hamiltonian (2.9) is essentially the same as for a free particle, the question arises whether the $osp(2, 1)$ likely extends into the super-Schrödinger algebra [8].

For further results the Reader is referred to [9].

References

- [1] L. C. Biedenharn, Phys. Rev. **126**, 845 (1962); L. C. Biedenharn and N. V. V. Swamy, Phys. Rev. **5B**, 1353 (1964); M. Berrondo and H. V. McIntosh, Journ. Math. Phys. **11**, 125 (1970).
- [2] G. Temple, *The General Principles of Quantum Mechanics*, New York: Methuen (1948); P. C. Martin and R. J. Glauber, Phys. Rev. **109**, 1307 (1958).
- [3] F. Bloore and P. A. Horváthy, Journ. Math. Phys. **33**, 1869 (1992).
- [4] E. D'Hoker and L. Vinet, Phys. Rev. Lett. **55**, 1043 (1986); E. D'Hoker, V. A. Kostelecky and L. Vinet, in *Dynamical Groups and Spectrum Generating Algebras*, p. 339-367, World Scientific, Singapore (1988).
- [5] E. D'Hoker and L. Vinet, Comm. Math. Phys. **97**, 391-427 (1985).
- [6] Y. Kazama, C. N. Yang and A. S. Goldhaber, Phys. Rev. **D15**, 2287 (1977); Y. Kazama and C. N. Yang, Phys. Rev. **D15**, 2300 (1977).
- [7] R. Jackiw, Ann. Phys. (N.Y.) **129**, 183 (1980).
- [8] J. P. Gauntlett, J. Gomis and P. K. Townsend, Phys. Lett. **248B**, 288 (1990); C. Duval and P. A. Horváthy, Journ. Math. Phys. **35**, 2516 (1994).
- [9] P. A. Horváthy, A. J. Macfarlane and J.-W. van Holten, Phys. Lett. **B 486**, 346-352 (2000).